

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>01 AUG 2015</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Hydrogen cyanide related deaths and detection in the blood</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) <b>Bebarta V. S.,</b>				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>UU</b>	18. NUMBER OF PAGES <b>2</b>	19a. NAME OF RESPONSIBLE PERSON
a REPORT <b>unclassified</b>	b ABSTRACT <b>unclassified</b>	c THIS PAGE <b>unclassified</b>			

## LETTER TO THE EDITOR

# Hydrogen cyanide related deaths and detection in the blood

Vikhyat S. Bebarta

*Department of Emergency Medicine, San Antonio Military Medical Center, San Antonio, TX, USA*

In response to manuscript: Stamyr K, Thelander G, Ernstgard L, Ahlner J, Johanson G. Swedish forensic data 1992–2009 suggest hydrogen cyanide as an important cause of death in fire victims. *Inhal Toxicol*. Feb 2012;24(3):194–199.

We read with interest the manuscript by Stamyr et al. which describes the cyanide levels of deceased patients exposed to structural fires [1]. This unique paper is important and adds more scientific support to what many clinicians feel is true, that cyanide is a common cause of toxicity and death after smoke inhalation. For many reasons, detection of cyanide toxicity is difficult, including accessibility of obtaining cyanide levels, ambiguity of cyanide related symptoms, and other reasons described by authors.

However, we felt two questions require clarification by the authors. The authors report that a spectrophotometric method was used to detect cyanide levels prior to 2002. However, many patients in Europe received hydroxocobalamin prehospital and in the hospital for presumed cyanide toxicity [2]. Hydroxocobalamin has light absorption characteristics that interfere with colorimetric and spectrophotometric laboratory measurements [3]. As examples, artificial elevations in creatinine, glucose, and bilirubin have been reported, as have decrease alanine aminotransferase, and unpredictable results for creatine phosphokinase, phosphate, and lactate dehydrogenase [4]. Urinalyses are often uninterruptable [3]. Thus, we are concerned that deceased patients in the database may have received hydroxocobalamin before death that could have caused spectrophotometric interference.

They authors did not state if hydroxocobalamin interferes with their spectrophotometric cyanide assay used for detection before 2002.

Secondly, hydroxocobalamin binds cyanide well. In particular, investigators have reported that once this antidote is infused in sufficient doses, cyanide levels are undetectable [5]. If patients in this study had received hydroxocobalamin, their cyanide levels may be reduced or undetectable as measured by either analytical method used for cyanide detection. The authors did not report which, if any, patients received an antidote, specifically hydroxocobalamin and thus potentially underestimating the proportion of patients that had detectable cyanide levels.

## References

1. Stamyr K, Thelander G, Ernstgard L, Ahlner J, Johanson G. 2012. Swedish forensic data 1992–2009 suggest hydrogen cyanide as an important cause of death in fire victims. *Inhal Toxicol* 24:194–199.
2. Borron SW, Baud FJ, Barriot P, Imbert M, Bismuth C. 2007. Prospective study of hydroxocobalamin for acute cyanide poisoning in smoke inhalation. *Ann Emerg Med* 49:794–801, e791–e792.
3. Beckerman N, Leikin SM, Aitchinson R, Yen M, Wills BK. 2009. Laboratory interferences with the newer cyanide antidote: hydroxocobalamin. *Semin Diagn Pathol* 26:49–52.
4. Curry SC, Connor DA, Raschke RA. 1994. Effect of the cyanide antidote hydroxocobalamin on commonly ordered serum chemistry studies. *Ann Emerg Med* 24:65–67.
5. Bebarta VS, Tanen DA, Laird J, Dixon PS, Valtier S, Bush A. 2010. Hydroxocobalamin and sodium thiosulfate versus sodium nitrite and sodium thiosulfate in the treatment of acute cyanide toxicity in a swine (*Sus scrofa*) model. *Ann Emerg Med* 55:345–351.

Copyright of Inhalation Toxicology is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.